

1. The intermolecular attractive forces would be greatest in which of the following molecules: CH_4 , CH_2Cl_2 and CO_2 .
 - (A) CH_4
 - (B) CH_2Cl_2
 - (C) CO_2
 - (D) the same in all three molecules
 - (E) no way to predict this property


2. The melting point of the halogens (Group VII: F_2 , Cl_2 , Br_2 , I_2) increases as you move down the column. This is because:
 - (A) Induced dipole effects increase with molecular weight.
 - (B) The smaller molecules have more hydrogen bonding.
 - (C) I_2 is a solid at room temperature.
 - (D) The larger molecules are more polar.
 - (E) both A and D above.

3. The predominant intermolecular attractive force in solid xenon is:
 - (A) ionic
 - (B) covalent
 - (C) metallic
 - (D) dipole-dipole
 - (E) induced dipole-induced dipole

4. Urea, $\text{O}=\text{C}(\text{NH}_2)_2$ would be predicted to be most soluble in which solvent:
(A) H_2O (B) CCl_4 (C) CCl_3H (D) CH_4 (E) C_3H_8

5. Calculate the energy, in kJ, required to convert a 36.00 g sample of solid water at 0°C to liquid water at 0°C . The molecular weight of water is 18.00 g/mol. For water, $\Delta H_{\text{vap}}=40.7 \text{ kJ/mol}$ and $\Delta H_{\text{fusion}}=6.0 \text{ kJ/mol}$.
(A) 6.0 (B) 12.0 (C) 20.4 (D) 40.7 (E) 81.4

Use the following phase diagram for a substance to answer questions 6 and 7.



6. A sample of the substance is placed in a container at a temperature of 85°C and the pressure is fixed at 10 mm Hg. Under these conditions what phase(s) exist?


- (A) solid only
- (B) liquid only
- (C) gas only
- (D) solid and liquid
- (E) liquid and gas

7. The substance is placed in a container at the same original conditions as in problem number 6. The substance is then cooled to 0 °C at *constant pressure*. What phase(s) will be present at the final conditions?

- (A) solid only
- (B) liquid only
- (C) gas only
- (D) solid in equilibrium with liquid
- (E) solid in equilibrium with gas

8. The heat of vaporization of water is 40.7 kJ/mol. It boils at 100 °C at 1 atm. At what temperature does water boil at the top of Mt. Washington, where the atmospheric pressure is 0.80 atm?

- (A) 89 °C
- (B) 94 °C
- (C) 97 °C
- (D) 99.5 °C
- (E) 105 °C




9. The unit cell of the mineral perovskite is shown above. The unit cell described by the Ca^{2+} ions is _____.

- (A) simple cubic
- (B) face centered cubic
- (C) body centered cubic
- (D) hexagonal close packing
- (E) tetrahedral

10. The formula of perovskite is _____.

- (A) CaTiO_3
- (B) CaTiO_6
- (C) Ca_2TiO_3
- (D) Ca_2TiO_6
- (E) Ca_8TiO_6

11. The decomposition of SO_2Cl_2 is a first-order reaction:



The rate constant is 0.17 1/sec. What is the rate of the reaction at an initial concentration $[\text{SO}_2\text{Cl}_2] = 1.2 \times 10^{-3} \text{ mol/L}$? The rate is in units of mol/(L sec).


- (A) 2.0×10^{-4} (B) 6.3×10^{-4} (C) 1.2×10^{-3} (D) 7.1×10^{-3} (E) 3.2×10^{-2}

12. The reaction

$2 \text{NO} + 2 \text{H}_2 \rightarrow \text{N}_2 + 2 \text{H}_2\text{O}$ is second order in $[\text{NO}]$ and first order in $[\text{H}_2]$
If the rate of the reaction is 0.55 M s^{-1} when $[\text{NO}] = 0.90 \text{ M}$ and $[\text{H}_2] = 0.45 \text{ M}$, what is the rate constant in $\text{M}^{-2}\text{s}^{-1}$?

- (A) 0.68 (B) 1.36 (C) 1.51 (D) 2.72 (E) 8.90

The following data were collected for the rate of disappearance of NH_3 in the reaction:




Experiment	$[\text{BF}_3](\text{M})$	$[\text{NH}_3](\text{M})$	Initial rate M/s
1	0.0126	0.0125	1.41×10^{-2}
2	0.0252	0.0250	1.13×10^{-1}
3	0.0252	0.0125	5.64×10^{-2}

13. What is the rate law for the reaction?

- (A) $\text{Rate} = k[\text{BF}_3][\text{NH}_3]$ (D) $\text{Rate} = k[\text{BF}_3]$
(B) $\text{Rate} = k[\text{BF}_3]^2[\text{NH}_3]$ (E) $\text{Rate} = k[\text{BF}_3]^2[\text{NH}_3]^2$
(C) $\text{Rate} = k[\text{BF}_3][\text{NH}_3]^2$

14. In a study of the gas phase decomposition of dimethyl ether at 500°C ,




the concentration of CH_3OCH_3 was followed as a function of time. It was found that a graph of $\ln[\text{CH}_3\text{OCH}_3]$ versus time in seconds gave a straight line with a slope of $-6.00 \times 10^{-4} \text{ s}^{-1}$ and a y-intercept of -1.99

Based on this plot, the reaction is _____ order in CH_3OCH_3
and the rate constant for the reaction is _____ s^{-1} .

- (A) zero, 6.00×10^{-4}
(B) first, 6.00×10^{-4}
(C) first, -7.42
(D) second, 6.00×10^{-4}
(E) second, -7.42

15. The reaction



was studied at 25°C. The following results were obtained where

$$\text{Rate} = - \frac{d[\text{S}_2\text{O}_8^{2-}]}{dt}$$

$[\text{I}^-]_0$ (mol/L)	$[\text{S}_2\text{O}_8^{2-}]_0$ (mol/L)	Initial rate (mol/L s)
0.080	0.040	12.50×10^{-6}
0.040	0.040	6.25×10^{-6}
0.080	0.020	6.25×10^{-6}
0.032	0.040	5.01×10^{-6}
0.060	0.030	7.02×10^{-6}

The rate equation for this reaction is:

(A) $R = k[\text{I}^-][\text{S}_2\text{O}_8^{2-}]$

(B) $R = k[\text{I}^-]^2[\text{S}_2\text{O}_8^{2-}]$

(C) $R = k[\text{I}^-][\text{S}_2\text{O}_8^{2-}]^2$

(D) $R = k[\text{I}^-]^2[\text{S}_2\text{O}_8^{2-}]^2$

(E) $R = k[\text{I}^-]$

The half life for the radioactive decay of ^{14}C is 5720 years.
16. What is the first order rate constant for this process?

(A) 5720 year

(B) $1.75 \times 10^{-4} \text{ year}^{-1}$

(C) $1.21 \times 10^{-4} \text{ year}^{-1}$


(D) $8.25 \times 10^3 \text{ year}^{-1}$

(E) not enough information

17. If a 1.000 g sample of ^{14}C were to sit for 3000 years, how much ^{14}C would remain?

- (A) 0.524 g (D) 0.363 g
(B) 0.784 g (E) 0.696 g
(C) 0.250 g

18. The decomposition of ammonia on a platinum surface at 856 °C




is zero order in NH_3 with a rate constant of $1.50 \times 10^{-6} \text{ Ms}^{-1}$.

If the initial concentration of NH_3 is 0.00116 M, the concentration of NH_3 will be 0.000213 M after _____ seconds have passed.

- (A) 142 (D) 1.02×10^6
(B) 631 (E) 1.13×10^6
(C) 773


19. The gas phase decomposition of nitrosyl chloride at 500 K



is second order in NOCl with a rate constant of $5.90 \times 10^{-2} \text{ M}^{-1}\text{s}^{-1}$.

If the initial concentration of NOCl is 0.080 M, the concentration of NOCl will be 0.020 M after _____ seconds have passed.

- (A) 0.34 (D) 636
(B) 1.02 (E) 847
(C) 212



20. A reaction profile for the reaction $2 O_2 \rightarrow O_3 + O$ is shown above. Which of the following is true (choose only one)
- (A) This reaction is endothermic, the value of the activation energy is 392 kJ, and ΔE is +411 kJ.
 - (B) This reaction is endothermic, the value of the activation energy is 803 kJ, and ΔE is +392 kJ.
 - (C) This reaction is endothermic, the value of the activation energy is 411 kJ, and ΔE is +392 kJ.
 - (D) This reaction is exothermic, the value of the activation energy is 411 kJ, and ΔE is -392 kJ.
 - (E) This reaction is exothermic, the value of the activation energy is 392 kJ, and ΔE is -411 kJ.